

ELLS FILE (1)31

OFFICE OF NAVAL RESEARCH

Contract N00014-87-K-0541

R & T Code 4133020

Technical Report No. 10

Modelling the In Situ Infrared Reflection-Absorption Spectra of the Diffuse Layer

by

P.W. Faguy, S. McCullough and W.R. Fawcett

Prepared for Presentation

at

The Electrochemical Society Meeting Los Angeles, CA, May 1989

May 1, 1989

Reproduction in whole or in part is permitted for any purpose of the United States Government

"This document has been approved for public release and sale; its distribution is unlimited"



REPORT DOCUMENTATION PAGE	1. REPORT NO.	2	1. Resigners A	etrision No.
	Situ Infrared Re e Diffuse Layer	flection-Absorpt	ion S. Report Date.	1, 1989
Faguy, S. M	cCullough and W.R	. Fawcett	& Performing C	rganization Root, Na.
Department of Ch			IQ. Project/Tee	b/Wart Unit No.
University of Ca Davis, CA 9561				er Grannico Na. -87-K-0541
Office of Naval			12. Type of Rec	ort & Period Covered
800 N. Quincy Arlington, VA 2	2217-5000		14.	
Prepared for Pre Los Angeles, CA,	sentation at: The May 1989	Electrochemical	Society Meeti	ing,
36. Abstract (Limits 200 warran)				
method is applied organic electrol	ption spectra of d to spectra obtains ytes on gold election of the decision of	ined in the doubtrodes. κ_{ev}	de layer regio	n for aprotic,
		·	•	
17. Document Analysis & Occurre	649 649			
å. Montflors/Open-Guing Terme				
I	•			
	•			
e COMATI Puld/Group				
6. COMATI FINA/Group IS. Availability Statement		Uncl	Class (This Resert) assified. Class (This Page)	23. Ion. of Pages

75-Word Abstract Form
Extended Abstract must be submitted with the 75-Word Abstract by December 1, 1966

Los Angeles, California—May 7-12, 1989

Submit to: The Electrochemical Society, Inc 19 South Main Street, Penningto	in. NJ 00634-2006			to be assigned by the Society
With a copy to the Organizing Cl	neurmen In Si	tu Techniq	ues for Electr	rochemistry
(Title of Symposium)	•			
sponsored by Physical	1 Electroche	mistry		
(Division/Group)				
itte of paper	<u>In situ</u> Infr	ared Refle	ction-Absorpti	on Spectra
of the Diffuse	e Double Lay	er.		
Authors (Underline name of author presenting pa	. P W	Facuv. S	McCullough. W.	R. Fawcett
luthors (Underline name of author presenting pa	iper.) <u> </u>	<u> </u>		
Depa	artment of C	hemistry		
lusiness affiliation and address			Davida	
	fornia	M1.1.M1.11.W9		(916) 752-4261
(State or Cou			(ZIP Code)	(Telephone No.)
	(Type abetract in t	his area double	eneced \	
Using a stratified med	, ,,			nterface and
Gouy-Chapman-Stern theory	of the doubl	e laver a	method to mode	l infrared
		~		
reflection-absorption spec	tra of the d	liffuse lay	er is develope	ed. This method
is applied to spectra obta	ined in the	double lay	er region for	aprotic, organic
electrolytes on gold elect	ro des.		a	
i			4	
į.	NTIS GRA&I			
	Unannounced	_		
	Justificati	on		-′.op ₃
	D-			PECTED
	Distribution	on/		
	Availabil			
'		and/or	 ,	
	Dist Spe	cial		
	 			
lo you require any audiovisual equipment?	20-1		Has the Informatio	n in this abstract been
36 mm (2 x 2 in.) elide projector			presented verbally or published?	, submitted for publication
Overhead projector			□ Yee □ N	
Specify other (subject to availability and cost)				a, please provide the
is a full length paper on this work to be	submitted for Box	dety Journal pul	blication?	O No
Papers presented before a Society technical ma	eling become the non	party of the Societ	y and may not be sublished	ed elecuhera without written
permission of the Society. Papers presented at S	polety technical mee	tiogs must be author have been been been been been been been be	and may not be published by a member or spon	na ensemble without writish isored by an active member.
ine	ert name of Society	member author or	sponsor	••••••

Modelling the In Situ Infrared Reflection-Absorption Spectra of the Diffuse Double Layer.

P.W. Faguy, S. McCallough, W.R. Fawcett

Department of Chemistry, University of California, Davis, CA 95616-0395

The sensitivity of in sime IRRAS techniques 1 such as EMIRS and SNIFTIRS to molecular structure at the electrode / electrolyte interface arises from the coupling of two physical phenomena: the local increase in the electric field strength for parallel polarized light reflect at the surface of the electrode 2 and the potential drop across the double layer. In the EMIRS or SNIFTIRS experiment there is enhanced sensitivity to species present within a few wavelengths of the IR radiation from the electrode surface and only to those species which change due to the applied electrochemical potential. As the diffuse layer is typically 10 - 100 Å for these experiments 3 and as the large electromagnetic field at the metal surface decays over much longer distances 2, 2 - 20 µm, the resultant in sime spectra will be a function of the product of these two effects integrated over the complete diffuse layer. Thus with such techniques the diffuse layer is probed to almost the same extent as is the electrode surface.

This paper presents a method for calculating the in situ infrared reflection-absorption spectra for species present in the diffuse layer based on a stratified media model of the electrode / electrolyte interface, Gouy-Chapman-Stern theory and the attenuated total reflection IR spectra of the electrolyte solutions.

In order to calculate relection-absorption spectra it is necessary to construct an optical model of the interface. Figure 1 is a schematic drawing showing an N layer system consisting of a non-absorbing semi-infinite initial phase, N-2 planar finite strata and an absorbing semi-infinite final phase. This approach assumes that the media are isotropic, homogeneous, linear and that the dielectric constant is independent of position within an individual layer. A.5 Calculations of optical parameters for the in sine experiment, based on similar models, have been reported. 6.7

In this treatment, the matrix formalism developed by Hansen⁵ for the N layer system will be apply to a stratified medium where the 3rd through N-2 components represent the diffuse layer as is shown in Figure 1. Each of these N-4 strata will have a dielectric constant and characteristic thickness associated with it, obtained from bulk solution and solvent values and GSC theory. The real and the imaginary components of the dielectric constants are calculated using ATR measurements of the bulk components and the appropriate Fresnel equations.^{8,9}

Modelled and measured spectra for gold electrodes, using alkali perchlorate saits in dimethylformamide and acetonitrile solutions will be presented.

Acknowledgements

We gratefully acknowledge the support of the Office of Naval Research, Washington.

References

- A. Bewick and S. Pons in "Advances in Infrared and Raman Spectroscopy" Vol. 12, R.J.H Clark and R.E. Hester Eds., Heyden: Philadelphia, 1985, Chapter 1.
- J.D.E. McIntyre in "Advances in Electrochemistry and Electrochemical Engineering", Vol. 9, R.H Muller Ed.; Wiley & Sons: New York, 1973, Chapter 2.
- A. J. Bard and L.R. Faulkner "Electrochemical Methods", Wiley & Sons: New York, 1980, p506.
- O.S. Heavens "Optical Properties of Thin Solid Films", Dover: New York, 1965. Chapter 4.
- 5) W.N. Hansen, J. Opt. Soc. Amer. 1968, 58, 380.
- 6) H. Seki, K. Kunimatsu, W.G. Golden, Appl. Spectrosc. 1985, 39, 437.
- 7) D.S. Bethune, A.C. Luntz, J.K. Sass and D.K. Roe, Surf. Sci 1988, 197, 44.
- 8) J.E. Bertie and H.Y. Eysel, Appl. Spectrosc. 1985, 39, 392.
- B.C. Crawford, T.G. Goplen and D. Swanson in "Advances in Infrared and Raman Spectroscopy", Vol. 4, R.J.H. Clark and R.E. Hesser Eds., Heyden: Philadelphia, 1978, Chapter 2.

IR window		n,
bulk electrolyte		- n̂, d,
diffuse layer	:	- :
compact layer		- -

Figure 1. Multiple layer optical model of the electrode / electrolyte interface.